



## **FOOD ALLERGY**

Asthmatic: 🗆 Yes	s □ No (increased risk	for severe reaction)	
Allergen(s):			
Mother:	M Home #: _	M Work #:	M Cell #:
		F Work #:	
		Relationship:	
SYMPTOMS OF AN  • MOUTH  • THROAT  • SKIN  • STOMACH  • LUNG  • HEART	ALLERGIC REACTION No. Itching &swelling of I Itching, tightness in the Hives, itchy rash, swelling Nausea, abdominal of the state of th	MAY INCLUDE ANY/ALL OF TI ips, tongue or mouth, mouth throat, hoarseness, cough relling of face and extremities cramps, vomiting, diarrhea repetitive cough, wheezing ssing out"	HESE: "feels hot"
		ent is give immediately.	
STAFF MEMBER	RS INSTRUCTED:	☐ Classroom Teacher(s)	☐ Special Area Teacher(s)
□А	dministration	☐ Support Staff	☐ Transportation Staff
TREATMENT:	Rinse contact area w	vith water if appropriate	·
		otoms	
Benadryl ordered:	☐ Yes ☐ No	GiveBenadryl p	er provider's order
Call school nurse	Call parent/quardian if off s	school grounds	
	Call parent/guardian if off s	_	
	-	school grounds. instructions:	
IF INGESTION OF EPINE Preferred hospital increased heart rathospital by ambula	ed:  Yes  No Special  R SUSPECTED INGESTICE  PHRINE IS ORDERED, Go  if transported:  les a 20 minute response w  te. This is a normal response  ance. A staff member shoul	_	SYMPTOMS ARE PRESENT A TELY AND CALL 911.  dent may feel dizzy or have an ine should be transported to the emergency room if the parent
IF INGESTION OF EPINE Preferred hospital Epinephrine providincreased heart rathospital by ambula guardian or emerge	ed:	on of Allergen occurs, sometimes of the second of the seco	dent may feel dizzy or have an ine should be transported to the emergency room if the parent other students is present.
Finephrine ordered  IF INGESTION OF EPINE  Preferred hospital increased heart rate hospital by ambula guardian or emerged  Transportation  Special instruction  Healthcare Provider:	ed:	DN OF ALLERGEN OCCURS, SIVE EPHINEPHRINE IMMEDIA  vindow. After epinephrine, a stu se. Students receiving epinephr d accompany the student to the t and adequate supervision for coole on bus	dent may feel dizzy or have an ine should be transported to the emergency room if the parent other students is present.
Finephrine ordered  IF INGESTION OF EPINE Preferred hospital increased heart rate hospital by ambulate guardian or emerged  Transportation Special instruction  Healthcare Provider: Written by:	ed:	DN OF ALLERGEN OCCURS, SIVE EPHINEPHRINE IMMEDIA  vindow. After epinephrine, a stu se. Students receiving epinephr d accompany the student to the t and adequate supervision for occur  Dle on bus	SYMPTOMS ARE PRESENT ATELY AND CALL 911.  dent may feel dizzy or have an ine should be transported to the emergency room if the parent other students is present.  ailable on bus   Does not ride but